

Anikó KOVÁCS<sup>1\*</sup>, Vivien NAGY<sup>1</sup>, Szabolcs HOMOLYA<sup>1</sup>, Katalin Badakné KERTI<sup>1</sup>

<sup>1</sup>Department of Grain and Industrial Plant Processing, Hungarian University of Agriculture and Life Sciences, Budapest Hungary  
Email : kovacs.anikog4@uni-mate.hu

## ABSTRACT

From year to year the food industry has to follow the new food trends, the consumers' new demands, the possible changes of the ingredients and the supply system. Likewise, in recent years in many products' marketing, palm oil free mark is a positive sign for the costumers. This is due to the assumed negative impact of the palm oil cultivation on the rain forests, animal species, environment and human health. To this new trend the food industry has to adapt. All palm fat using companies have two choices. One is to only buy this ingredient from sustainable sources, and to join the Roundtable on Sustainable Palm Oil (RSPO). The other option is to use other fats instead of palm fat. However, due to the unique fatty acid composition and the several possible usages of palm fat, the substitution will be difficult, and the opportunities have to be investigated by product categories. In confectionary applications, one opportunity could be the mixing of fully hydrogenated fat with oil or with lauric fat. However, the suitability of the replacement needs to be examined from many perspectives, like rheological, textural, thermal behaviour and from sensory points of view.

Therefore, the goal of our research was to investigate the thermal and texture behaviour of palm fat and two different fat mixtures, as possible palm fat replacers.

## MATERIALS & METHODS

For substitution, we used fully hydrogenated rapeseed oil mixed in one case with sunflower oil and in the other case with coconut fat.

The mixing ratios were determined based on previous experiences.

Table 1: Investigated samples

	Samples
1	Palm fat - control
2	65% sunflower oil 35% fully hydrogenated rapeseed oil
3	95% coconut fat 5% fully hydrogenated rapeseed oil

We followed the thermal behaviour with differential scanning calorimetry (NETZSCH 3500) and the texture behaviour with penetrometry using conical pressure head at 25 °C (Stable Micro System TA-XT 2i).



Figure 1: NETZSCH DSC 3500



Figure 2: Stable Micro System TA-XT 2i texture analyser

## CONCLUSIONS

In conclusion, the coconut fat sample showed very promising values in case of the texture parameters. Also in the case of the thermal behaviour this samples crystallization peak temperature were closer to the palm fat. Further analyses are needed to decide the coconut fat mixture sample suitability as palm oil replacement, especially looking at its shelf life.

## RESULTS

According to our results (Figure 3.), the three fats' DSC curves showed different heating and cooling behaviours due to the different fat acid and triacyl glycerol compositions of the samples. However, for the crystallization behaviour all samples showed two peaks, and the coconut sample peak temperatures were close to the palm fat's.




Figure 3: Average DSC curves of the samples (n=3)



Figure 4: Average Hardness values of the samples (n=5)



Figure 5: Average Work of penetration values of the samples (n=5)



Figure 6: Average Adhesiveness values of the samples (n=5)

In case of texture measurement, we recorded the hardness (Figure 4.), work of penetration (Figure 5.) and adhesiveness values (Figure 6.). The results showed that the coconut sample values were very similar to the palm fat values in every parameter. There were no significant differences between these two samples. In case of the investigated texture parameters the coconut fat samples could be a substitution the control sample. However, the sunflower sample showed higher values in all cases than the palm fat, as it shown in the Figures.

## ACKNOWLEDGEMENTS

The research was supported by the European Union through the EFOP-3.6.3-VEKOP-16-2017-00005 project and by the Doctoral School of Food Science.

## REFERENCES

- N. Hinrichsen (2016)- Commercially available alternatives to palm oil, Lipid Technology Vol. 28, No. 3–4
- R. Cazzolla Gatti et al (2019)- Sustainable palm oil may not be so sustainable, Science of the Total Environment 652 (2019) 48-51

## KEYWORDS

substitution, palm fat, DSC, texture analyses